

On 4-dimensional cosmological models locally embedded in a 5-dimensional Ricci-flat space

A. G. Agnese* and M. La Camera†

*Dipartimento di Fisica dell'Università di Genova
Istituto Nazionale di Fisica Nucleare, Sezione di Genova
Via Dodecaneso 33, 16146 Genova, Italy*

Abstract

We employ a theorem due to Campbell to build some simple 4-dimensional cosmological models which originate from solutions describing waves propagating along the extra-dimension of a 5-dimensional Ricci-flat space. The dimensional reduction is performed in the Jordan frame according to the induced-matter theory of Wesson.

PACS numbers: 04.50.+h , 11.10.Kk

*Email: agnese@ge.infn.it

†Email: lacamera@ge.infn.it

An interesting version of 5-dimensional General Relativity has been developed in recent years by Wesson [1,2,3]. The central thesis of his induced-matter theory is that the 4D Einstein's field equations with matter

$$G_{\alpha\beta} = 8\pi T_{\alpha\beta} \tag{1}$$

are a subset of the 5D field equations for vacuum in terms of the Ricci tensor

$$R_{AB} = 0 \tag{2}$$

The theory also allows to obtain flat cosmological solutions containing the usual 4D perfect fluid energy-momentum tensor [4].

Due to its primary significance in the present context, it is worth quoting a theorem due to Campbell [5], which states as follows:

Theorem: *Any analytic Riemannian space $V_n(s, t)$ can be locally embedded in a Ricci-flat Riemannian space $V_{n+1}(s + 1, t)$ or $V_{n+1}(s, t + 1)$.*

This theorem has been recently brought to light by Romero et al. [6,7] and employed both in applying Wesson's method [1] and in investigating the embedding of lower-dimensional spacetimes .

In this Brief Note we consider the cosmological solution describing waves propagating in the extra-dimension of a (4+1)-dimensional Ricci-flat space and, after selecting particular modes, we obtain the corresponding (3+1)-dimensional cosmologies. The final result will be written in the Jordan frame,

according to the dimensional reduction prescribed by the “induced-matter theory of Wesson”.

We start from the 5D line element

$$ds_5^2 = e^\omega (dr^2 + d\Omega^2) - e^\nu dt^2 + e^\mu dl^2 \quad (3)$$

where $d\Omega^2 = d\vartheta^2 + \sin^2\vartheta d\varphi^2$ and l is the extra coordinate.

The metric coefficients ω , ν and μ will depend in general on both t and l , the dependence on r being ruled out in the absence of sources. The case when sources are present and the metric coefficients depend only on r , has been treated in a more general context in ref. [8]. Moreover we can put, by symmetry considerations, $\nu = \mu$.

The relevant Ricci equations are:

$$3\nu'\omega' - 3\omega'^2 - 2\nu'' - 6\omega'' + 3\dot{\nu}\dot{\omega} + 2\ddot{\nu} = 0 \quad (4a)$$

$$3\nu'\omega' + 2\nu'' + 3\dot{\nu}\dot{\omega} - 3\dot{\omega}^2 - 2\ddot{\nu} - 6\ddot{\omega} = 0 \quad (4b)$$

$$\omega'\dot{\nu} + \nu'\dot{\omega} - \omega'\dot{\omega} - 2\dot{\omega}' = 0 \quad (4c)$$

$$3\omega'^2 + 2\omega'' - 3\dot{\omega}^2 - 2\ddot{\omega} = 0 \quad (4d)$$

Here partial derivatives with respect to t and l are denoted by an overdot and a prime respectively.

One can immediately see that equation (4d) admits 3-brane wave solutions, propagating along the fifth dimension of a 5-dimensional bulk, of the form

$\omega_{\pm} = \omega(t \pm l)$ and $\nu_{\pm} = \nu(t \pm l)$. Denoting by an asterisk derivatives with respect to $t \pm l$, equations (4a), (4b) and (4c) all become, after substitution:

$$2 \nu_{\pm}^* \omega_{\pm}^* - \omega_{\pm}^{*2} - 2 \omega_{\pm}^{**} = 0 \quad (5)$$

Therefore, selecting a particular form of $\omega_{\pm}$, the other metric coefficients are given by

$$e^{\nu_{\pm}} = e^{\mu_{\pm}} = L \omega_{\pm}^* e^{\frac{\omega_{\pm}}{2}} \quad (6)$$

where L is a suitable constant of integration.

Starting from the above solutions, which describe waves propagating along the extra dimension of a 5-dimensional Ricci-flat spacetime with line element

$$ds_5^2 = e^{\omega_{\pm}} (dr^2 + r^2 d\Omega^2) + L \omega_{\pm}^* e^{\frac{\omega_{\pm}}{2}} (-dt^2 + dl^2) \quad (7)$$

we build some simple cosmological models in a 4-dimensional spacetime with line element

$$ds_4^2 = e^{\omega(t)} (dr^2 + r^2 d\Omega^2) - L \dot{\omega}(t) e^{\frac{\omega(t)}{2}} dt^2 \quad (8)$$

obtained from (7) by a section with a hypersurface at constant l chosen, without loss of generality, as $l = 0$.

Components of the Einstein tensor in mixed form, derived from the above metric (8), are

$$\begin{aligned} G_r^r = G_{\vartheta}^{\vartheta} = G_{\varphi}^{\varphi} &= - \frac{e^{-\frac{\omega}{2}} (\dot{\omega}^2 + \ddot{\omega})}{2 L \dot{\omega}} \\ G_t^t &= - \frac{3 e^{-\frac{\omega}{2}} \dot{\omega}}{4 L} \end{aligned} \quad (9)$$

We wish to match the terms in (9) with the components of the usual 4D perfect fluid energy-momentum tensor $T_{\alpha\beta} = (p + \rho) u_\alpha u_\beta + p g_{\alpha\beta}$. In our case the pressure and density are given by $T_r^r = p$ and $T_t^t = -\rho$, and therefore we can simply identify G_r^r with $8\pi p$ and G_t^t with $-8\pi\rho$.

Of course the choice of the function $\omega(t)$ is to a large extent arbitrary so we suggest, to make some physically meaningful examples, the following one:

$$\omega(t) = \alpha \ln \left(1 + \frac{t}{\alpha L} \right) \quad (10)$$

where α is an assignable constant.

As a consequence, the line element (8) becomes

$$ds_4^2 = \left(1 + \frac{t}{\alpha L} \right)^\alpha (dr^2 + r^2 d\Omega^2) - \left(1 + \frac{t}{\alpha L} \right)^{\frac{\alpha}{2}-1} dt^2 \quad (11)$$

and clearly characterizes a conformally flat spacetime when $\alpha + 2 = 0$.

To go further, it is useful to make in (11) the change of variable

$$\tau = \begin{cases} \frac{4\alpha L}{\alpha + 2} \left[\left(1 + \frac{t}{\alpha L} \right)^{\frac{\alpha+2}{4}} - 1 \right] & \text{if } \alpha + 2 \neq 0 \\ -2L \ln \left(1 - \frac{t}{2L} \right) & \text{if } \alpha + 2 = 0 \end{cases} \quad (12)$$

thus obtaining

$$ds_4^2 = \left(1 + \frac{(\alpha + 2)\tau}{4\alpha L} \right)^{\frac{4\alpha}{\alpha+2}} (dr^2 + r^2 d\Omega^2) - d\tau^2 \quad \text{if } \alpha + 2 \neq 0 \quad (13)$$

and

$$ds_4^2 = e^{\frac{\tau}{L}} (dr^2 + r^2 d\Omega^2) - d\tau^2 \quad \text{if } \alpha + 2 = 0 \quad (14)$$

where in both cases $1/(2L)$ represents the Hubble constant H_0 . Accordingly, pressure and density of the perfect fluid can be rewritten respectively as

$$8\pi p = \frac{\frac{2(1-\alpha)H_0^2}{\alpha}}{\left(1 + \frac{(\alpha+2)H_0\tau}{2\alpha}\right)^2} \quad 8\pi\rho = \frac{3H_0^2}{\left(1 + \frac{(\alpha+2)H_0\tau}{2\alpha}\right)^2} \quad (15)$$

and

$$8\pi p = -3H_0^2 \quad 8\pi\rho = 3H_0^2 \quad (16)$$

It is apparent that the case $\alpha + 2 = 0$ describes a de-Sitter Universe with cosmological constant $\Lambda = 3H_0^2$. On the other hand, the case $\alpha + 2 \neq 0$, provides the equation of state of radiation $3p = \rho$ when $\alpha = 2/3$, and the equation of state of matter $p = 0$ when $\alpha = 1$.

References

- [1] Wesson P. S., *Phys. Lett. B*, **276** (1992) 299.
- [2] Overduin J. M. and Wesson P. S., *Phys. Reports*, **283** (1997) 304.
- [3] Wesson P. S., *Space - Time - Matter. Modern Kaluza-Klein theory*,
(World Scientific, Singapore) 1999.
- [4] Billyard A. and Wesson P. S., *Gen. Rel. Grav.*, **28** (1996) 129.
- [5] Campbell J. E., *A Course of differential Geometry*, (Clarendon Press,
Oxford) 1926.
- [6] Romero C., Tavakol R. and Zalaletdinov R., *Gen. Rel. Grav.*, **28** (1996)
365.
- [7] Lidsey J. E., Romero C., Tavakol R. and Rippl S., gr-qc 9907040,
Preprint 1999.
- [8] Agnese A. G. and La Camera M., *Phys. Rev. D*, **58** (1998) 087504.